

www.BrundageGroup.com

Sepsis-3 SOFA Made Easy

Please include the word <u>severe</u> whenever diagnosing your patient using the 2016 Sepsis-3 SOFA criteria.

- The Society of Critical Care Medicine supports the Sepsis-3 (SOFA) scoring system to diagnose severe sepsis.
- The 2016 JAMA paper suggests that all sepsis is severe sepsis when using the SOFA criteria.

The patient must have an acute change in physical exam and/or laboratory findings consequent to infection to meet the criteria. Remember the diagnosis must be Present On Admission (POA) to be included in CMS (Medicare's) calculated expected mortality.

How sick is your patient, doctor? Are you documenting this effectively?

Sepsis-3 SOFA Score: Sequential Organ Failure Assessment			
	Acute change ≥ 2 points due to the infectious process supports sepsis		
System	0	1	2
Neurologic GCS	15	13-14	10-12
Respiratory PaO ₂ /FiO ₂ room air PaO ₂ , O ₂ sat	≥ 400 84, 95%	< 400 84, 95%	< 300 63, 91%
Cardiovascular	MAP ≥ 70 mmHg	MAP < 70 mmHg	Dopamine < 5 or Dobutamine (any)
Hepatic Bilirubin, mg/dL	< 1.2	1.2-1.9	2.0-5.9
Coagulation Platelets, x 1,000	≥ 150	< 150	< 100
Renal Creatinine, mg/dL	< 1.2	1.2-1.9	2.0-3.4
UOP, ml/d	≥ 500		< 500 (3 points)

Sepsis-3 definition: Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3), *JAMA*, 315(8);801-810, February 22, 2016 https://jamanetwork.com/journals/jama/fullarticle/2492881

Abbreviations:

PaO₂: partial pressure of oxygen; FiO₂: fraction of inspired oxygen;

MAP: Mean arterial pressure: [(SBP - DBP) / 0.33] + DBP

Catecholamine doses are in

μg/kg/min for at least 1 hour.