

Common Inpatient CC and MCCs

<u>CCs</u> (moderate severity)	<u>MCCs</u> (maximal severity)
Chronic systolic heart failure	Acute systolic HF
Chronic diastolic heart failure	Acute diastolic HF
Chronic respiratory failure	Acute respiratory failure
Mild/moderate malnutrition	Severe malnutrition
TIA	Stroke, acute
Acute renal failure/Acute kidney injury	Acute renal failure due to ATN (Remember you can use "possible, probable, likely, suspected" verbiage. You don't need a biopsy, just clinical intuition)
CKD stage 4 & 5	ESRD
Diabetic (type 1 or 2) gangrene	DM type 1 with DKA
	DM type 1 or 2 with coma (Think AMS with DM and hypo or hyperglycemia)
Unspecified shock	Specified shock (i.e. septic, cardiogenic, etc.)
Bacteremia	Sepsis
COPD with acute exacerbation	Pneumonia
Demand ischemia	NSTEMI or STEMI
Dementia with behavior disturbances	Metabolic/toxic encephalopathy
	Stage 3 or 4 pressure ulcer (That is present on admission--> POA)
Persistent atrial fib	
Acute blood loss anemia	
Hemiplegia or hemiparesis	
Morbid obesity with alveolar hypoventilation	