



www.BrundageGroup.com

Outpatient HCCs

ICD-10 Codes to Screen for HCC Relative Weight	HCC RW
Diabetes Mellitus Often DM is coded without complication (even when patients have manifestations of their DM) <ul style="list-style-type: none"> • Uncomplicated DM has only 1/3 the HCC relative weight of DM with <u>any</u> complication • Consider Diabetes (type 1 or 2) with hyper- or hypoglycemia* 	 0.106 0.307
Pneumonia HCC relative weight is available with specificity of describing pneumonia <ul style="list-style-type: none"> • Unspecified Pneumonia, CAP, HAP, HCAP <i>all without HCC RW</i>) • Lobar – Localize with a stethoscope or a chest X-ray • Aspiration 	 0.000 0.164 0.612
Respiratory <ul style="list-style-type: none"> • Home Oxygen Use • Chronic respiratory failure - Supplemental oxygen use⁺ 	 0.000 0.314
Bronchitis <ul style="list-style-type: none"> • Acute bronchitis has no HCC RW • Chronic bronchitis has the same RW as COPD 	 0.000 0.335
Alcohol Use Alcohol use, abuse and dependency <ul style="list-style-type: none"> • Alcohol abuse • Alcohol dependency has HCC weight • Alcohol dependency in remission (“on the wagon” in recovery) is HCC RW for life 	 0.000 0.368 0.368

*When clinically supported

⁺ Requirement of long-term supplemental oxygen, not transient or situational